

## 9 Myths and Misunderstandings about Menopausal Hormone Therapy (MHT)

The International Menopause Society is grateful for permission to adapt for global use the information sheet that was originally produced by the Australasian Menopause Society. The medical and scientific information provided might not be relevant to a particular woman's circumstances and should always be discussed with her own health-care provider.

www.menopause.org.au



# 9 Myths and Misunderstandings about Menopausal Hormone Therapy (MHT)

www.menopauseinfo.org



**Menopausal Hormonal Therapy (MHT)** is the best treatment to improve hot flushes and quality of life for menopausal women.

In the early 2000s, the Women's Health Initiative (WHI) studies of postmenopausal women caused confusion among women, the media and doctors.

New information about MHT and the WHI studies mean doctors better understand the risks and benefits of MHT.

## **Main Points**:

- Many of the myths about MHT come from the WHI studies of 2002 and 2004. New information about MHT means **doctors better understand the risks and benefits** of MHT.
- MHT is the **best way to control menopausal symptoms** and gives some women health benefits.
- Combined MHT (oestrogen plus progestogen) or oestrogen alone cause **no significant increase in breast cancer or risk of heart disease in women aged 50 to 59 or in women who start treatment within 10 years of menopause**. New information about MHT mean doctors better understand the risks and benefits of MHT.

## 1. MHT will make you put on weight

Women who use MHT **do not gain any more weight** than women who do not use MHT. Ageing, social, lifestyle and medical factors are the main causes of midlife weight gain. The hormonal changes of menopause do cause fat to move from the hips to the abdomen.

## MYTH

## 2. Breast cancer is the most common cause of death in postmenopausal women

**Fear of breast cancer is a major reason why women do not use MHT**. Many postmenopausal women believe they are more likely to die from breast cancer than heart disease or stroke. The opposite is true. For example, in Australia in 2014, 12 in every 100 women died from coronary heart disease, nine in 100 from stroke and four in 100 from breast cancer.

## MISUNDERSTANDING

## 3. One quarter of women who take MHT get breast cancer

This misunderstanding came about because some journalists **incorrectly** reported the early WHI study results. New information combined with better understanding suggests that:

• Five years taking combined MHT (oestrogen plus progestogen) did **not increase breast cancer risk** in women aged 50 to 59 or in women who started treatment within 10 years of menopause. After 13 years, women had a small increase in breast cancer risk (nine extra cases of breast cancer per 10,000 women)

• Seven years taking oestrogen only MHT did **not increase breast cancer risk** in women aged 50 to 59 or in women who started treatment within 10 years of menopause. After 13 years, there was still no increased risk of breast cancer.

#### MISUNDERSTANDING

## 4. MHT increases the risk of heart disease

An analysis of all studies (40,410 women) showed **MHT did not increase the number of deaths from heart and blood vessel disease or heart attacks**. MHT also did not increase the number of cases of angina in healthy women or in women with pre-existing heart and blood vessel disease.

#### MYTH

## 5. A blood test is necessary to diagnose menopause

Blood tests for hormone levels, and other tests, are **not needed to diagnose menopause**. A woman is considered postmenopausal when she is over the age of 45 and has had at least 12 months without a period. Blood tests can be helpful for women who are younger than 40 years of age or who have had a hysterectomy and have menopausal symptoms.

#### MYTH

## 6. Complementary medicines and therapies are as effective as MHT and are safer

**MHT remains the most effective way to control menopausal symptoms**. Some complementary medicines and therapies are promoted as natural and safe without evidence that they work. Some products (such as soy) should be avoided if you are unable to take prescribed MHT for safety reasons. Often there is no way to know if complementary therapies are safe or uncontaminated, especially if they are bought online. Speak with your doctor about complementary medicines or therapies. They might not be suitable for your situation. If you have any concerns or questions about options to manage your menopausal symptoms, visit your doctor.

#### MYTH

## 7. Compounded bioidentical hormone therapy is safer than conventional MHT

Doctors **advise against** the use of compounded bioidentical hormone therapy. "Bioidentical hormones" are chemically the same as those produced by the body. Some MHTs prescribed by your doctor are "bioidentical". There is no evidence that compounded bioidentical hormone therapy is better than prescribed MHT. Compounded bioidentical hormone therapy offers no advantages and many disadvantages because there are:

- no regulations for their use
- no standards for quality of manufacturing
- no testing of the products for negative effects, quality or safety
- serious side effects such as endometrial cancer
- possible higher cost

## MISUNDERSTANDING


### 8. All progestogens have the same risks

Therapy combining progestogens and oestrogen is used to reduce the risk of cancer of the uterus in women who still have a uterus. Progestogen is a term that covers both progesterone (naturally occurring in humans) and progestins (synthetic progesterones). **Different types of progestogens have different risks**. Your doctor can discuss the different MHT options available and work with you to reduce your risk.

## MISUNDERSTANDING

## 9. Nonhormonal medications are as effective as MHT for hot flushes

Evidence suggests **nonhormonal treatments are not as effective as MHT**, although more studies are needed. You should ask your doctor about nonhormonal treatment options if you are not able to use MHT for medical reasons or you do not want to use MHT.





1